

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/353121629>

ANTI-OXIDANT ACTIVITY OF AQUEOUS AND ORGANIC EXTRACTS FROM KENYAN RUELLIA PROSTRATA

Article · January 2017

CITATIONS

3

READS

81

6 authors, including:



Christine Wangia

Jomo Kenyatta University of Agriculture and Technology

11 PUBLICATIONS 30 CITATIONS

SEE PROFILE



Jennifer Orwa

Kenya Medical Research Institute

56 PUBLICATIONS 1,146 CITATIONS

SEE PROFILE



Francis W Muregi

Mount Kenya University

38 PUBLICATIONS 1,354 CITATIONS

SEE PROFILE



Cheruiyot Kipyegon

Jomo Kenyatta University of Agriculture and Technology

22 PUBLICATIONS 200 CITATIONS

SEE PROFILE



Received on 25 August, 2016; received in revised form, 15 November, 2016; accepted, 21 November, 2016; published 01 March, 2017

ANTI-OXIDANT ACTIVITY OF AQUEOUS AND ORGANIC EXTRACTS FROM KENYAN *RUPELLIA PROSTRATA*

Christine O. Wangia^{*1}, Jennifer A. Orwa², Francis W. Muregi³, Patrick G. Kareru⁴, Kipyegon Cheruiyot⁴ and Japheth Kibet²

Department of Pharmacology¹, Department of Chemistry⁴, Jomo Kenyatta University of Agriculture and Technology, Nairobi, Kenya

Centre for Traditional Medicine and Drug Research², Kenya Medical Research Institute, P.O. Box 54840-00200 Nairobi, Kenya

Department of Biological Sciences³, Mount Kenya University, P.O. Box 342-01000, Thika, Kenya

Keywords:

Ruellia prostrata, Antioxidant, Phytochemicals, Flavonoids, Tannins, Saponins

Correspondence to Author:

Christine O. Wangia


Department of Pharmacology,
Jomo Kenyatta University of
Agriculture and Technology, P.O.
Box 62000-00200, Nairobi, Kenya

E-mail: cwangia@jkuat.ac.ke

ABSTRACT: *Ruellia* species belong to Acanthaceae family, and are perennial creepers with widespread medicinal uses including analgesic and anti-inflammatory activity. The objective of this study was to determine the qualitative phytochemical constituents and anti-oxidant activity of crude extracts from whole plant parts of Kenyan *Ruellia* species viz. *Ruellia prostrata*. Whole parts of the plants were collected, air-dried under shade and organic extraction was done by cold maceration using ethyl acetate and methanol. Aqueous extraction was done by boiling. Anti-oxidant activity was performed based on the ability of the aqueous, methanolic and ethyl acetate extracts to scavenge free radicals produced by 2,2-Diphenyl-1-picrylhydrazyl (DPPH). Ascorbic acid was used as the standard. Anti-oxidant tests were done at eight concentrations (3.9, 7.8, 15.6, 31.3, 62.5, 125, 250 and 500 µg/ml). The anti-oxidant activity of both plant extracts and ascorbic acid increased with increase in concentration. Methanolic and ethyl acetate extracts exhibited a higher anti-oxidant activity with IC₅₀ values of 20.58 and 22.26 µg/ml respectively, relative to aqueous extract (IC₅₀ of 51.92µg/ml). Ascorbic acid standard exhibited high anti-oxidant activity with IC₅₀ value of 2.11µg/ml. The phytochemical screening tests were based on visual observation of colour change and precipitate formation. Phytochemical analysis revealed the presence of terpenoids, saponins, flavonoids, tannins and cardiac glycosides. Flavonoids and tannins are a major group of compounds that act as primary anti-oxidants. The presence of these compounds could attribute to the potent anti-oxidant activity of *R. prostrata* extracts. From this study, it was concluded that *Ruellia prostrata* could have anti-inflammatory activity.

INTRODUCTION: Medicinal plants play a critical role in the health sector of ancient and modern cultures. They are widely used as alternative therapeutic tools for prevention and treatment of many diseases¹.

Despite modern screening and discovery of drugs, plant-based medicines remain the lead compounds to the discovery of valuable drugs². Drugs of plant origin are locally available, often cheaper, with minimal side effects^{3, 4}. Nowadays, herbal drugs form an integral part of alternative medicine⁵. Plants have been found to produce and accumulate secondary metabolites which *Ruellia prostrata* (Acanthaceae) is a perennial fall into various groups based on their biosynthetic origin. Humans have used these plant secondary metabolites as medicines and flavoring agents⁶.

QUICK RESPONSE CODE 	DOI: 10.13040/IJPSR.0975-8232.8(3).1282-86
	Article can be accessed online on: www.ijpsr.com
DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.8(3).1282-86	

Some plant secondary metabolites belong to the glycoside group, flavonoids, tannins, saponins and alkaloids. Some of these compounds have biological effects including anti-oxidant activity⁷. Normal biochemical reactions in the body produce several free radicals that play a critical role in the development of various diseases such as asthma, dementia, cancer, arthritis and Parkinson's disease^{8, 9}. Anti-oxidants from plant materials play a significant role in termination of these free radicals, thereby protecting the body from these diseases. Herb mainly known for its traditional use as an anti-inflammatory and anti-cancer against the epidermis of nasopharynx region and possesses wound healing properties^{10, 11}.

Arthritis is an inflammatory disorder which manifests by destructive arthropathy and extra-articular manifestations, leading to severe disability and premature mortality¹². The present study aimed at screening the phytoconstituents and anti-oxidant potential of Kenyan *Ruellia prostrata* to make this plant useful for formulation of analgesic and anti-arthritic drug for management of arthritis.

MATERIALS AND METHODS:

Chemicals: Ascorbic acid, 2,2-diphenyl-1-picrylhydrazyl (DPPH), ethyl acetate and methanol were of analytical grade.

Sample collection and preparation: Whole plant parts of *R. prostrata* were collected from Mwala sub-County, Machakos County, in December, 2010, taxonomised and the Voucher specimen (Ref. no. UoN/2010/598 December 16, 2010) deposited at the Department of Botany Herbarium, University of Nairobi. The plant sample was washed in water, chopped and shade-dried for three weeks. Once completely dried, grinding into coarse powder was done using an electrical grinder (manufactured by Jomo Kenyatta University of Agriculture and Technology, Mechanical Engineering Department) and extracted with methanol, ethyl-acetate and distilled water. The extracts were prepared by soaking 50g of plant powder in 500ml of (Methanol, ethyl-acetate and distilled water) for 72 hours. The extracts from organic solvents were concentrated using a rotary evaporator (BUCHI R-200) at 45°C. The aqueous extract from distilled water was dried in a freeze-drier (Christ Alpha 1-4 LD). All the extracts were stored at 4°C until

required for phytochemical screening and antioxidant activity.

Phytochemical analysis: Qualitative phytochemical screening was performed as described by¹³. The extracts were tested for saponins, alkaloids, phenolic compounds, flavonoids, phytosterols and glycosides. The screening tests were based on observation of colour change and precipitate formation.

Test for alkaloids: Mayer's test: In this test, 1mL of Mayer's reagent (potassium mercuric iodine) was added to 1mL of the test solution and observed for a white precipitate, which is a positive indicator for the presence of alkaloids.

Test for tannins: Ferric chloride test: In this test, 0.5mL of 5% ferric chloride solution was added to 0.5mL of the sample solution. Formation of a dark-green colour would indicate the presence of tannins.

Test for flavonoids: To 1 mL of crude extract, 3 drops of ammonia solution were added followed by 0.5mL of concentrated hydrochloric acid. The formation of pale brown colouration would indicate the presence of flavonoids.

Test for glycosides (Keller-Killian test): Glycosides were tested by adding 1mL of 3.5% ferric chloride in acetic acid to 1mL of the sample solution followed by careful drop-wise addition of 1.5mL concentrated sulphuric acid by the sides of the test tube to form a separate layer at the bottom. A brown ring at the interface due to the presence of de-oxy sugar would be a characteristic of cardenolides and a pale green colour in the upper layer due to the steroidal nucleus would be taken as the indicator for the presence of cardiac glycosides.

Test for saponins: Plant extract (0.5 g) was added into 5 ml of distilled water. The mixture was shaken vigorously and observed for the formation of foam that would persist for 15 minutes. Development of a stable emulsion on addition of three drops of olive oil to the formed foam would confirm the presence of saponins¹⁴.

Test for terpenoids: Crude extract (0.5g) was dissolved in 5mL of chloroform and evaporated to dryness. To this, 2mL of concentrated sulphuric

acid was added and heated for about 2 minutes. A reddish brown colouration at the interface would indicate the presence of terpenoids.

2, 2-Diphenyl - 1- Picrylhydrazyl (DPPH) Scavenging Activity:

The DPPH scavenging activity on the plant samples was done according to the procedure¹⁵. DPPH (0.5 mM) solution in methanol (3.94 mg in 100 ml) was freshly prepared. Concentrations ranging from 3.9-500 µg/ml for both extracts and standards were prepared. Equal volume of DPPH solution was added. The samples were kept in a dark place and after 30 minutes the absorbance was measured at 517 nm. The percentage of the DPPH radical scavenging was calculated using the equation given below:

Percentage (%) Inhibition =

$$\frac{(\text{Absorbance of}_{\text{DPPH}} - \text{absorbance of}_{\text{SAMPLE}}) \times 100}{\text{Absorbance}_{\text{DPPH}}}$$

Determination of inhibitory concentration (IC₅₀):

The results for anti-oxidant activity were presented as a line graph for three experiments. The concentration required to inhibit 50% of DPPH radical (IC₅₀) was determined using the regression line of probit according to the log₁₀ of the extract concentration.

RESULTS AND DISCUSSION: Phytochemical screening of aqueous, ethyl acetate and methanolic extracts of *Ruellia prostrata* aerial parts as shown in **Table 1**.

TABLE 1: PHYTOCHEMICAL SCREENING OF AQUEOUS, ETHYL ACETATE AND METHANOLIC EXTRACTS OF RUELLIA PROSTRATA AERIAL PARTS

Phytochemical	Type of solvents		
	Methanol	Aqueous	Ethyl acetate
Saponins	+	+	-
Phenolics	+	+	-
Alkaloids	-	-	-
Glycosides	+	+	+
Flavonoids	+	+	+
Terpenoids	+	+	+

Plants have an almost limitless ability to synthesize substances most of which are secondary metabolites of which at least 12,000 compounds have been isolated¹⁶. Phytochemical substances play an important role in plant defence against

microorganism, stress as well as interspecies protections. These plant components have been used as drugs for many years. Thus, phytochemical screening serves as the initial step in predicting the types of potential active compounds from plants¹⁷. Phytochemical screening of *Ruellia prostrata* aerial parts tested positive for saponins, phenolics, flavonoids, glycosides and terpenoids. Ethyl acetate extract showed absence of saponins and phenolics whereas all the extracts showed absence of alkaloids. Among these compounds, flavonoids, saponins and phenolics are important secondary metabolites and may be compounds responsible for medicinal values of the plant.

Antioxidant results for aqueous, methanolic and ethyl-acetate extracts of *Ruellia prostrata* species are presented in **Fig. 1**.




FIG. 1: RESULTS SHOWING THE ANTI-OXIDANT ACTIVITY OF AQUEOUS, METHANOLIC AND ETHYL-ACETATE EXTRACTS OF RUELLIA PROSTRATA SPECIES

Analysis of anti-oxidant activity data using t-test revealed that there was statistically significant difference in anti-oxidant activities for methanol, aqueous and ethyl acetate extract ($p < 0.05$). This could be attributed to the difference in proportions of the active components that are responsible for anti-oxidant activity. The difference in anti-oxidant activity could also be due to the difference in solubility of individual phytochemicals in solvents of extraction. Ascorbic acid in comparison with the activity of the three extracts showed significant difference in activity ($p < 0.05$).

However, ascorbic acid exhibited 99% inhibition at 15.6 µg/ml. The higher extract concentration resulted in higher anti-oxidant inhibition compared with lower concentration suggesting a concentration-dependent response.

This is indicated by the fact that increasing the concentration of the plant extracts increased the proportion of the chemical ingredient with pharmaceutical value in the plant extract¹⁸.

The IC₅₀ values of *Ruellia prostrata* aqueous, methanol and ethyl acetate extracts required to inhibit 50% of DPPH activity are presented in **Table 2**.

TABLE 2: IC₅₀ VALUES (UG/ML) FOR ANTI-OXIDANT ACTIVITY OF AQUEOUS, METHANOLIC AND ETHYL-ACETATE EXTRACTS OF *RUELLIA PROSTRATA* SPECIES

	Type of extract	IC ₅₀ values (ug/ml)	Lower Bound	Upper Bound
1.	Aqueous extract	51.92	28.28	100.78
2.	Methanolic extract	20.58	5.23	49.17
3.	Ethyl-acetate extract	22.26	5.97	52.46
4	Ascorbic acid	2.11	0.00	6.16

In the present study, the DPPH (1, 1- Diphenyl- 2-picrylhydrazyl) assay was conducted on aqueous, methanolic and ethyl-acetate extracts of *Ruellia prostrata*. Methanolic extract had the lowest IC₅₀ value (20.58 µg/mL), followed by ethyl-acetate extract (22.26 µg/mL) and aqueous extract (51.92 µg/mL). These extracts showed a significant DPPH scavenging activity when compared with positive control (ascorbic acid) which had an IC₅₀ value of 2.11µg/mL. DPPH has a stable centred-nitrogen free radical which can be easily and effectively scavenged by anti-oxidants.

This free radical accepts an electron or hydrogen radical to become a stable diamagnetic molecule. The anti-oxidant activity occurs when anti-oxidant molecules react with the radicals resulting to scavenging of the radical by hydrogen donation¹⁹. Phenolics and flavonoids play a key role in anti-oxidant activity since they are highly effective electron donors²⁰. These compounds are documented to possess anti-carcinogenic and anti-inflammatory properties²¹. Flavonoids are polyphenolic compounds that play a fundamental role in stabilizing lipid oxidation and are associated with anti-oxidant activity²². As described by²³, phenolic compounds may have an inhibitory consequence on mutagenesis and carcinogenesis in humans when as much as 1.0 g is ingested daily from diet rich in fruits and vegetables.

CONCLUSION: *Ruellia prostrata* species contain flavonoids and phenolic compounds which act as primary anti-oxidants or free- radical scavengers. The presence of these compounds could be attributed to the potent anti-oxidant activity useful for formulation of analgesic and anti-arthritis preparations.

ACKNOWLEDGEMENTS: The authors are thankful to Kenya Medical Research Institute and the Department of Chemistry (JKUAT) for technical support. Special thanks also to Research, Production and Extension (RPE) Division of Jomo Kenyatta University of Agriculture and Technology for funding the project.

REFERENCES:

1. Nagavani V and Rao TR: Evaluation of antioxidant potential and qualitative analysis of major polyphenols by RP-HPLC in *Nymphaea nouchali* Burm flowers. International Journal of Pharmacy and Pharmaceutical Sciences 2004; 2: 98-104.
2. Buenz F, Scheneppe D and Motley TJ: Technique: bio prospecting historical herbal texts by hunting for new leads in old tomes. Trends in pharmacological Sciences 2004; 25: 494-498.
3. Park E, Rhee H, Jung H, Ju S, Hong S, Yang H and Kim, K: Anti-inflammatory effects of a combined herbal preparation (RAH13) of *Phellodendron amurense* and *Coptis chinensis* in animal models of inflammation. Phytotherapy research 2007; 746-750.
4. Raj BA, Murugamani V, Madhuri B. Preliminary phytochemical investigation of *Givotia moluccana* Stem. Int J Res Pharm Biomed Sci. 2011; 2(3): 1307-1313.
5. Park EJ and Pezzutto JM: Botanicals in Cancer Chemo preventives. Cancer and Metastasis Reviews 2002; 21:231-255
6. Samuni-Blank M, Izhaki I, Dearing M, Gerchman Y, Trabelcy B, Lotan A and Arad Z : Intraspecific directed deterrence by the mustard oil bomb in a desert plant. Current biology 2012; 1-3.
7. Packer L, Rimbach G and Virgili F: Antioxidant activity and biologic properties of a procyanidin-rich extract from pine (*Pinus maritima*) bark, pycnogenol. Free Radical.Biology. Medicine 1999; 27: 704-724.
8. Lai LS and Chou ST: Studies on the antioxidative activities of Hsiantiao (*Mesona procumbens*.Hemsl.) leaf gum. Journal of Agriculture 2001; 49: 963-968.
9. Uy, MM, Garcia KI. Evaluation of the antioxidant properties of the leaf extracts of Philippine medicinal plants *Casuarina equisetifolia* Linn, *Cyperus brevifolius* (Rottb) Hassk, *Drymoglossum piloselloides* Linn, *Ixora chinensis* Lam, and *Piper abbreviatum* Opiz. AAB Bioflux. 2015; 7(2):71-79.

10. Jain SC and Jain R: Ethnobotanical Survey of Sariska and Siliserh Regions from Alwar District of Rajasthan, India. Ethnobotanical Leaflets 2009; 13:171-88.
11. Patel M, Verma R, Srivastav P. Antioxidant Activity of *Boerhavia diffusa* Extract. International Journal of Pharmacognosy and Phytochemical Research. 2014; 6: 598 – 605.
12. Aletaha D, Neogi T, Silman A, Funovits J, Felson D and Bingham C: Rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. Arthritis Rheumatoid 2010; 62(9):2569-2581.
13. Harbone J: Phytochemical methods, a guide to modern techniques of plant analysis, 3rd Edition, Chapman and Hall International Edition, New York 1998.
14. Trease G and Evans W: Pharmacognosy 2nd Edition. Braille Tiridel and Macmillan Publishers 1989.
15. Bouhlali EDT, Ramchoun M, Alem C, Ghafoor K, Ennassir J, Zegzouti YF. Functional composition and antioxidant activities of eight Moroccan date fruit varieties (*Phoenix dactylifera L.*). Journal of the Saudi Society of Agricultural Sciences. 2015.
16. Ghoshal S, Prasad B and Lakshmi V: Antiamoebic activity of *Piper longum* fruits against *Entamoeba histolytica* *in vitro* and *in vivo*. Journal of Ethnopharmacology 2006; 50:167-170.
17. Chew YL, Chan EW, Tan, PL, Lim YY, Stanslas J and Goh JK: Assessment of phytochemical content, polyphenolic composition, antioxidant and antibacterial Activities of Leguminosae medicinal plants in Peninsular Malaysia. BMC Complementary and Alternative Medicine 2011; 11:12.
18. Debella A: Manual for phytochemical screening of medicinal plants. Ethiopian health and Nutrition Research Institute, Addis Ababa, Ethiopia 2002.
19. Rackova L, Oblozinsky M, Kostalova D, Kettmann V and Bezakova L: Free radical scavenging activity and lipoxygenase inhibition of *Mahonia aquifolium* extract and isoquinoline alkaloids. Journal of Inflammation 2007; 4:15.
20. Van-Acker S, Van-Balen G, Vanden-Berg D, Bast A and Vander-Vijgh S: Influence of iron chelation on the antioxidant activity of flavonoids. Biochemistry. Pharmacology 1998; 56: 935-943.
21. Wong C, Cheng H and Chen F: A systematic survey of antioxidant activity of 30 Chinese medicinal plants using the ferric reducing antioxidant power assay. Food Chemistry 2006; 97: 705-711.
22. Yen GC, Duh P and Tsai CL: Relationship between antioxidant activity and maturity of peanut hulls. Journal of Agriculture. Food Chemistry 1993; 41: 67-70.
23. Tanaka M, Kuei W, Nagashima Y and Taguchi T: Application of antioxidative maillard reaction products from histidine and glucose to sardine products. Nippon Suisan Gakkai Shil 1998; 54:1409-1414.

How to cite this article:

Wangia CO, Orwa JA, Muregi FW, Kareru PG, Cheruiyot K and Kibet J: Anti-oxidant activity of aqueous and organic extracts from kenyan *ruellia prostrata*. Int J Pharm Sci Res 2017; 8(3): 1282-86.doi: 10.13040/IJPSR.0975-8232.8(3).1282-86.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)